

SECTION VII. THE NORTHERN AND NORTHWESTERN EDGE OF THE BOHEMIAN MASS

Introduction

This work treats the Lausitz, the Elbe valley region, the North Bohemian Tertiary area, the Erzgebirge, the Vogtland, and the Fichtel Mountains. This whole region forms the northern and northwestern edge of the Bohemian Mass, and thus follows the discussion of the Silesian ore region.

The discussion is preceded by a review of the over-all structure of the Bohemian Mass, which is followed by a geological-tectonic review of the area.

The deposits in the area under discussion are described in detail. The area is subdivided into:

- A. Pre-Varistian deposits
- B. Deposits of the Varistian orogenesis
- C. Saxonian deposits

The pre-Varistian deposits are of no great importance. Most of them are syngenetic pyrite deposits situated in pre-Upper Devonian rocks. Most of them have been subjected to Varistian effects. The intensity of the Varistian influence increases from northwest to southeast, with local metamatoses being caused by intruded granites.

The edge zone is dominated by deposits of the Varistian orogenesis. The great diversity allows a new insight into the mutual effect of magma, tectonics, and deposits.

The following types of deposits are discussed:

- 1. Pre- and syntectonic ore deposits of predominantly contact-metamorphic genesis
- 2. Contact-metascmatic and contact-metamorphous ore deposits, bound to post-tectonic granites
- 3. ¹⁰⁰Pneumatolytic and pneumatolytic-catathermal deposits with tin and tungsten ores
- 4. Hydrothermal vein fillings

The temporal, spatial, and causal relations between the deposits, magma, and tectonics are discussed in conjunction with each individual area. The discussion of the dependencies, found to occur within small areas, is supplemented by the corresponding discussions, covering broad areas of the units of higher order.

The Saxonian deposits are of no economic importance. However, they supply excellent evidence on the geological position of the more recent mineralization. Only one real, so to speak, autochthonous Saxonian deposit can be detected, while most of the fissure fillings are mobilized. Their position depends, on the one hand, on Varistian deposits in the sub-base, and, on the other hand, on Tertiary thermal springs.

The last section summarizes the results and emphasizes the relations between the geological and structural factors and the ore deposits of the northern and northwestern edge zone of the Bohemian Mass. It shows dependencies and mutual effects between small units, e.g., the individual late-Varistian granite intrusions bearing the Varistian mineralization, as well as relationships covering a large area, which can be observed throughout the entire northwestern, northern, and northeastern peripheral zone. A deep zone of weakness is marked by the late Varistian granites and by the Tertiary magmatism. This zone of weakness was used by the most recent late Varistian granites, which contain many volatile components, for their uprise, so that the deposits connected with them are bound to this zone.

Simultaneously, findings on the distribution of the Varistian deposits within the entire Bohemian Mass were developed.

The real primary deposit formation in connection with the Varistian orogenesis ends with the Rotliegendes.

It can be proved that most of the fissure fillings are mobilized in the Cretaceous and in the Tertiary. Only the lead-zinc ore deposit of the Bohemian Mountains near Ronstok on the Elbe is bound to the Tertiary magmatism and connected with a center of eruption.

Table of Contents

- I. Geological Summary
 - A. Over-all structure of the Bohemian Mass
 - B. The northern and northwestern edge zone of the Bohemian Mass
 - 1. The area of the Lausitz granite
 - 2. The Erzgebirge, with the shale system of the Elbe valley and the Vogtland
 - 3. The Fichtel Mountains
- II. The deposit units in the northern and northwestern edge zone of the Bohemian Mass
 - A. Pre-Varistian deposits in the northern and northwestern edge zone of the Bohemian Mass
 - 1. General summary of the syngenetic and contact-metamorphous ore deposits
 - 2. Syngenetic pyrite deposits in pre-Upper Devonian rocks
 - a) Syngenetic pyrite deposits, with only slight or no metamorphous influence
 - 1) Pyrite in the East Thuringian shale mountains and in the Franconian Forest
 - 2) Pyrite in the alum shales of northwestern Bohemia
 - b) Syngenetic pyrite deposits in pre-Upper Devonian rock, with Varistian influence
 - 1) Klingenthal - Grasnitz in the Erzgebirge
 - 2) Johanngeorgenstadt
 - 3) Kupferberg, Neufang, and Adlerhuetten in Upper Franconia
 - 4) Sparneck in the Fichtel Mountains
 - 5) Pyrite deposits near Waldsassen - Tirschenreuth - Baernau
 - 6) The pyrite deposits of the katasone of the Bavarian Forest
 - Bodenmais
 - Lam in the Bavarian Forest

3. The geological-tectonic position of the pyrite deposits
in the western part of the Bohemian Mass
- B. The deposits of the Varistian orogenesis
 1. Varistian tectonic phases and Varistian magmatism in the
northern and northwestern edge of the Bohemian Mass
 2. Pre- and syntectonic ore deposits, of predominantly contact-
metamorphic genesis
 3. Contact-metasomatic and contact-metamorphous ore deposits
bound to post-tectonic granites
 - a) Schwarzenberg - Breitenbrunn - Rittersgruen
 - b) Geyer - Thum - Ehrenfriedersdorf
 - c) Goldenhoehe, northwest of Gottesgab
 - d) The deposits between Aue and Bockau
 - e) Other ore deposits in the western Erzgebirge
 - f) Berggiesshuelbel in the shale system of the Elbe valley
 4. The magmatic and tectonic dependencies of the Varistian
contact deposits
 5. Pneumatolytic and pneumatolytic-catathermic deposits with
tin and tungsten ores
 - a) The tin-tungsten ore region of the eastern Erzgebirge
 - 1) Graupen
 - 2) Zeidelweide near Fuerstenau
 - 3) Zinnwald
 - 4) Altenburg in the eastern Erzgebirge
 - 5) Raupennest and Paradies-Fundgrube
 - 6) Badisdorf
 - 7) Hoher Busch
 - 8) Lauenstein - Loewenhain
 - 9) Hegelschoebe, west of Baerenstein
 - 10) Frauenberg near Frauendorf
 - b) Magmatic and tectonic dependencies of the tin and tin-
tungsten deposits of the eastern Erzgebirge

- c) The granites of the western Erzgebirge and their development
- d) The tin, tin-tungsten and tungsten ore deposits, bound to the granite of Eibenstock - Neudeck and its peripheral associates
 - 1) "Hoffnung Gottes" mine on the Kaffberg near Rittersgruen
 - 2) Ochsendracht tin veins, east northeast of Johann-georgenstadt
 - 3) Waidmann shaft, north of Zwittermuehl
 - 4) Henneberg, southwest of Johanngeorgenstadt
 - 5) Oberjugel, southwest of Johanngeorgenstadt
 - 6) "Gabe Gottes" mine at Doenitzgrund
 - 7) "Rote Grube" mine at Doenitzgrund
 - 8) Hirschenstand
 - 9) Fruehbuss
 - 10) Hengstererben
 - 11) Platten
 - 12) Baeringen
 - 13) Tannenberg near Tannenbergsthal
 - 14) Gottesberg, southeast of Tannenbergsthal
 - 15) Rothau
 - 16) Montanus and Helene mines near Schneeberg
 - 17) Zachorlau near Schneeberg
 - 18) Martin Roemer mine, east of Kirchberg
 - 19) Pechtelsgruen near Lengenfeld
 - 20) Stangengruen, east of Lengenfeld
 - 21) Tirpersdorf near Oelsnitz in the Vogtland
 - 22) Pillmannsgruen
 - 23) Oelsnitz in the Vogtland
- e) The tin and tin-tungsten deposits bound to the Kaiserwald granite

- 1) Schlaggenwald - Schoenfeld
- 2) Lauterbach
- f) The tin ore deposits bound to the granite knobs of Ehrenfriedersdorf-Ceyer
- g) The tin ore deposits bound to the denuded granite anticline of Schlettau in the ore vein region of Annaberg
- h) The tin ore deposits of Pobershau, Marienberg and Seiffen, not bound to immediately detectable granites
- i) The dependence of the tin and tungsten deposits of the western Erzgebirge on the magma and tectonics
- j) The tin ore deposit region of the granite of the Fichtel Mountains
 - 1) Tin ore in the Schneeberg massif near Weissenstadt
 - 2) Tin ore near Wernerereuth
- k) The zinc-tungsten ore deposits in the edge zone of the Bohemian Mass and their dependence on tectonics and magma
6. Hydrothermal vein fillings
 - a) Summary and subdivision
 - b) The hydrothermal deposits in the Erzgebirge
 - 1) The veins in the region of the Freiberg gneiss knob
 - (1) The Freiberg region
 - (2) The Scharfenberg region near Meissen
 - 2) Hydrothermal deposits in the region of the eastern Erzgebirge tin-bearing granites
 - (1) Tellnitz
 - (2) Sachsenhoehe near Lauenstein
 - (3) Badisdorf
 - 3) Hydrothermal deposits bound to the Eibenstock granite and its peripheral associates
 - (1) Schneeberg and Oberschlema

- (2) Johanngeorgenstadt
- (3) Joachimsthal
- (4) Gottesgab and Niederschlag
- 4) The veins of Annaberg
- 5) The veins of Marienberg
- 6) Smaller individual regions
 - (1) Klostergrab
 - (2) Katharinaberg
- 7) Hydrothermal deposits in the region of the Kaiserwald granite
 - (1) Schoenficht
 - (2) Duerrmaul near Marienbad
- 8) The spatial, temporal and causal relationships between the Varistian deposits, magma and tectonics in the Erzgebirge
- c) The hydrothermal ore veins of the Vogtland
 - 1) Cata- to mesothermal mineralization above the deep-lying granite ridge of Bergen - Eichigt
 - 2) Meso- to telethermal fissure fillings near Schoenbrunn, Boesenbrunn, Wiedersberg, and Jocketa
 - 3) Telethermal lead ore veins at Bleistadt
 - 4) Telethermal mercury ore veins at Hartenstein and Schoenbach
 - 5) Magmatic and tectonic dependencies of the hydrothermal deposits in the Vogtland
- d) Hydrothermal deposits in the Fichtel Mountains
 - 1) Specularite veins near Fichtelberg
 - 2) The hydrothermal gold quartz veins of Brandholz in the Fichtel Mountains
 - 3) The hydrothermal lead-zinc ore veins of Erbendorf in the Upper Palatinate
 - 4) Magmatic and tectonic dependencies of the hydrothermal deposits of the Fichtel Mountains

e) Hydrothermal ore deposits in the region of the Lausitz granites

- 1) Hydrothermal nickeliferous pyrrhotite deposits at Schland on the Spree River and at Schweidrich near Schluckenau
- 2) Hydrothermal lead-zinc ore deposit at St. Georgenthal
- 3) Other ores in the Lausitz granite itself
- 4) The significance of the lack of deposits in regard to the geological position of the Lausitz granite region

C. Saxonian deposits

- 1) Hydrothermal, Tertiary mobilized mineralization in the region of the marginal fault of the Erzgebirge
- 2) Hydrothermal ore deposit of Rongstock as a consequence of the Tertiary volcanism in the North Bohemian basin
- 3) The genetic importance of the Rongstock mineralization

III. The relationship between the geological and structural factors and the ore deposits of the northern and northwestern edge zone of the Bohemian Mass

Illustrations in the Text

- Fig. 1:** The lower ore stratum and the phycoidal quartzites in the Vogtland-Thuringian Ordovician
- Fig. 2:** The chamositic iron ores of the Lower Silurian in the Vogtland-Thuringian Upper Tremadoc
- Fig. 3:** Schematic drawing of the Kupferberg deposit
- Fig. 4:** Geological conditions in the region of the pyrite deposit of Sparneck in the Fichtel Mountains
- Fig. 5:** Graphic representation of the age sequence of the minerals of the ore deposits of Joehstadt, Pressnitz and Kupferberg
- Fig. 6:** Graphic representation of the intensity of the most important minerals of the ore deposits of Joehstadt, Pressnitz and Kupferberg
- Fig. 7:** Graphic representation of the age sequence of the minerals of the ore deposits of Schwarzenberg - Breitenbrunn
- Fig. 8:** Graphic representation of the intensity of the most important minerals of the ore deposits of Schwarzenberg - Breitenbrunn
- Fig. 9:** Graphic representation of the age sequence of the mineralization of the ore deposits of Berggiesshuel
- Fig. 10:** Aplitic slabby inserts in the Eibenstock granite
- Fig. 11:** Map of the granite massif zone of the western Erzgebirge
- Fig. 12:** Schematic representation of the tin ore body of the Tannenberg mine
- Fig. 13:** The wolframite content of the shearing fissures of Zachorlen, bound to a definite, clearly limited pressure area
- Fig. 14:** Schematic representation of the mineralization of the tin ore deposit of Ehrenfriedersdorf
- Fig. 15:** H/F (Huebnerite/Ferberite) coefficients of various wolframite deposits
- Fig. 16:** Isothermal areas with the corresponding deposit zones in the region of the Sadisdorf tin-bearing granites

- Fig. 17: Zonary mineralization in the Schneeberg region
- Fig. 18: The specularite quartz veins near Fichtelberg in the Fichtel Mountains
- Fig. 19: Schematic profile through the gold quartz veins of Brandholz and the specularite quartz veins of Fichtelberg in the Fichtel Mountains
- Fig. 20: Cenomanian sandstone containing fluorite and quartz, located above quartz porphyry and below Turonian clay marl
- Fig. 21: Geological map of the Rongstock essexite and its surroundings
- Fig. 22: Rongstock essexite with hydrothermal mineralization of veins and fissures
- Fig. 23: Simplified, geological-tectonic map of the Fichtel Mountains and the Franconian Forest

Illustrations in Appendix

- Fig. 1: General map of the core of the Bohemian Mass and the northern edge zones
- Fig. 2: Geological map of the northern and northwestern edge zones of the Bohemian Mass
- Fig. 3: Deposits of uranium ores and of radioactive springs above 100 Maché units in the region of the Bohemian Mass
- Fig. 4: Zone of weakness on the northern edge of the Bohemian Mass
- Fig. 5: The pyrite and contact deposits in the west and northwest of the Bohemian Mass
- Fig. 6: Cross-section of the pyrite deposit of Graslitz - Klingenthal
- Fig. 7: The ore Deposits of the western Erzgebirge
- Fig. 8: Map of the ore deposits of the Erzgebirge, the Vogtland, and the shale system of the Elbe valley
- Fig. 9: The tin and tin-tungsten deposits of the eastern Erzgebirge
- Fig. 10: The Zinnwald granite massif in the eastern Erzgebirge
- Fig. 11: The Sadisdorf tin deposits

Fig. 12: The tin and tungsten ore deposits of the Eibenstock massif
and its peripheral associates

Fig. 13: Kohlgrub mine, Hirschenstand

Fig. 14: The ore deposits of the Eibenstock granite and its peripheral
associates

Fig. 15: The tin- and tin-tungsten deposits of Schoenfeld - Schlaggenwald

Fig. 16: Map of the veins in the eastern Erzgebirge

Fig. 17: Map of the veins of Annaberg and Marienberg

Fig. 18: Map of the veins of the western Erzgebirge

Fig. 19: Post-Cretaceous fluorite-barite mineralization in the region of
the Tertiary marginal fault of the Erzgebirge, between Teplitz
and Tetschen

Fig. 20: The mineralized fissures of the Vogtland with epithermal to
telethermal fissure fillings

(NOTE: The following map, although not listed in the Table of Contents,
apparently belongs to this volume: The ore deposits of the Erzgebirge.)